



DOH COVID-19 Vaccine Schedule

COVID-19 vaccination (minimum age: 6 months [Moderna Spikevax], 5 years [Pfizer-BioNTech Comirnaty], 12 years [Novavax Nuvaxovid, Moderna mNEXSPIKE])

Routine Vaccine Schedule for Healthy Persons Ages 6 Months-18 Years

[AAP Immunization Schedule](#)

Age 6-23 months	
Vaccination Status	Recommended Doses
Unvaccinated	<ul style="list-style-type: none">2 doses Moderna at 0, 4–8 weeks
Incomplete initial vaccination series with: <ul style="list-style-type: none">1 dose Moderna or 1 or more doses Pfizer-BioNTech	<ul style="list-style-type: none">Complete initial series with 1 dose Moderna 4–8 weeks after most recent dose
Completed initial vaccination series	<ul style="list-style-type: none">1 dose Moderna at least 8 weeks after the most recent dose

Special situations

Age 2-18 years in the following risk groups:

- Persons at high risk of severe COVID-19
- Residents of long-term care facilities or other congregate settings
- Persons who have never been vaccinated against COVID-19
- Persons whose household contacts are at high risk for severe COVID-19

Age 2–11 years

- 1 dose of Moderna or Pfizer-BioNTech (**for those 5 years and older**) regardless of previous vaccination status at least 8 weeks after the most recent dose

Age 12-18 years

- 1 dose of Moderna, Pfizer-BioNTech, or Novavax regardless of previous vaccination status at least 8 weeks after the most recent dose
- 1 dose of Moderna mNEXSPIKE for people who have previously received a COVID-19 vaccine. This dose should be administered at least 12 weeks after the last dose was received

Children 2 through 18 years of age not included in the risk groups above whose parent or guardian desires their protection from COVID-19 should be offered a single dose of age-appropriate COVID-19 vaccine.

Routine Vaccine Schedule for Healthy Adults 19 years and Older

[Adult Immunization Schedule | AAFP](#)

Age 19-64 years	
Vaccination Status	Recommended Doses
Unvaccinated	<ul style="list-style-type: none">• 1 dose Moderna or Pfizer-BioNTech• 2 doses Novavax at 0, 3–8 weeks
Previously vaccinated with: <ul style="list-style-type: none">• 1 or more doses Moderna or Pfizer-BioNTech• 1 dose Novavax• 2 or more doses Novavax• 1 or more doses Janssen	<ul style="list-style-type: none">• 1 dose Moderna or Novavax or Pfizer-BioNTech at least 8 weeks after the most recent dose• 1 dose Novavax 3–8 weeks after most recent dose. If more than 8 weeks after most recent dose, administer 1 dose Moderna or Novavax or Pfizer-BioNTech• 1 dose Moderna or Novavax or Pfizer-BioNTech at least 8 weeks after the most recent dose• 1 dose Moderna or Novavax or Pfizer-BioNTech

Age 65 years and older	
Vaccination Status	Recommended Doses
Unvaccinated	<ul style="list-style-type: none"> • 1 dose Moderna or Pfizer-BioNTech OR • 2 doses Novavax at 0, 3–8 weeks AND • Administer additional dose of Moderna or Novavax or Pfizer-BioNTech 6 months later (minimum interval 2 months) for persons 65 and older
Previously vaccinated with: <ul style="list-style-type: none"> • 1 or more doses Moderna or Pfizer-BioNTech • 1 dose Novavax • 2 or more doses Novavax • 1 or more doses Janssen 	Administer additional dose of Moderna or Novavax or Pfizer-BioNTech 6 months later (minimum interval 2 months) for persons 65 and older in addition to the doses recommended below. <ul style="list-style-type: none"> • 1 dose Moderna or Novavax or Pfizer-BioNTech at least 8 weeks after the most recent dose • 1 dose Novavax 3–8 weeks after most recent dose. If more than 8 weeks after most recent dose, administer 1 dose Moderna or Novavax or Pfizer-BioNTech • 1 dose Moderna or Novavax or Pfizer-BioNTech at least 8 weeks after the most recent dose • 1 dose Moderna or Novavax or Pfizer-BioNTech

Vaccine Schedule for *Immunocompromised Persons 6 Months and Older*

[AAP Immunization Schedule](#)

[Adult Immunization Schedule | AAFP](#)

Age 6 months-4 years	
Vaccination Status	Recommended Doses
Unvaccinated	<ul style="list-style-type: none"> • 4 doses (3-dose initial series Moderna at 0, 4 weeks, and at least 4 weeks after dose 2, followed by 1 dose Moderna 6 months later [minimum interval 2 months])
Previously vaccinated with:	

<ul style="list-style-type: none"> 1 dose Moderna 2 doses Moderna or 1 or more doses Pfizer-BioNTech 	<ul style="list-style-type: none"> Complete initial series with 2 doses Moderna at least 4 weeks apart (administer dose 1 4 weeks after most recent dose), followed by 1 dose Moderna 6 months later (minimum interval 2 months) Complete initial series with 1 dose Moderna at least 4 weeks after most recent dose, followed by 1 dose Moderna 6 months later (minimum interval 2 months)
Completed initial 3-dose vaccination series with: <ul style="list-style-type: none"> 3 or more doses Moderna or Pfizer-BioNTech 	<ul style="list-style-type: none"> 2 doses Moderna 6 months apart (minimum interval 2 months). Administer dose 1 at least 8 weeks after the most recent dose

Age 5-11 years	
Vaccination Status	Recommended Doses
Unvaccinated	<ul style="list-style-type: none"> 4 doses (3-dose initial series Moderna at 0, 4 weeks, and at least 4 weeks after dose 2, followed by 1 dose Moderna or Pfizer-BioNTech 6 months later [minimum interval 2 months]) 4 doses (3-dose initial series Pfizer-BioNTech at 0, 3 weeks, and at least 4 weeks after dose 2, followed by 1 dose Moderna or Pfizer-BioNTech 6 months later [minimum interval 2 months])
Incomplete initial 3-dose vaccination series with: <ul style="list-style-type: none"> 1 dose Moderna 2 doses Moderna 	<ul style="list-style-type: none"> Complete initial series with 2 doses Moderna at least 4 weeks apart (administer dose 1 4 weeks after most recent dose), followed by 1 dose Moderna or Pfizer-BioNTech 6 months later (minimum interval 2 months) Complete initial series with 1 dose Moderna at least 4 weeks after most recent dose, followed by 1 dose Moderna or

<ul style="list-style-type: none"> 1 dose Pfizer-BioNTech 2 doses Pfizer-BioNTech 	<p>PfizerBioNTech 6 months later (minimum interval 2 months)</p> <ul style="list-style-type: none"> Complete initial series with 2 doses Pfizer-BioNTech at least 4 weeks apart (administer dose 1 3 weeks after most recent dose), followed by 1 dose Moderna or PfizerBioNTech 6 months later (minimum interval 2 months) Complete initial series with 1 dose Pfizer-BioNTech at least 4 weeks after most recent dose, followed by 1 dose Moderna or Pfizer-BioNTech 6 months later (minimum interval 2 months)
<p>Completed initial 3-dose vaccination series with:</p> <ul style="list-style-type: none"> 3 or more doses Moderna or 3 or more doses PfizerBioNTech 	<p>2 doses Moderna or Pfizer-BioNTech 6 months apart (minimum interval 2 months). Administer dose 1 at least 8 weeks after the most recent dose</p>

Age 12-18 years	
Vaccination Status	Recommended Doses
Unvaccinated	<ul style="list-style-type: none"> 4 doses (3-dose initial series Moderna at 0, 4 weeks, and at least 4 weeks after dose 2, followed by 1 dose Moderna or Novavax or Pfizer-BioNTech 6 months later [minimum interval 2 months]) 4 doses (3-dose initial series Pfizer-BioNTech at 0, 3 weeks, and at least 4 weeks after dose 2, followed by 1 dose Moderna or Novavax or Pfizer-BioNTech 6 months later [minimum interval 2 months]) 3 doses (2-dose initial series Novavax at 0, 3 weeks, followed by 1 dose Moderna or Novavax or Pfizer-BioNTech 6 months later [minimum interval 2 months])
<p>Incomplete initial vaccination series with:</p> <ul style="list-style-type: none"> 1 dose Moderna 	

<ul style="list-style-type: none"> • 2 doses Moderna • 1 dose Pfizer-BioNTech • 2 doses Pfizer-BioNTech • 1 dose Novavax 	<ul style="list-style-type: none"> • Complete initial series with 2 doses Moderna at least 4 weeks apart (administer dose 1 4 weeks after most recent dose), followed by 1 dose Moderna or Novavax or Pfizer-BioNTech 6 months later (minimum interval 2 months) • Complete initial series with 1 dose Moderna at least 4 weeks after most recent dose, followed by 1 dose Moderna or Novavax or PfizerBioNTech 6 months later (minimum interval 2 months) • Complete initial series with 2 doses Pfizer-BioNTech at least 4 weeks apart (administer dose 1 3 weeks after most recent dose), followed by 1 dose Moderna or Novavax or PfizerBioNTech 6 months later (minimum interval 2 months) • Complete initial series with 1 dose Pfizer-BioNTech at least 4 weeks after most recent dose, followed by 1 dose Moderna or Novavax or Pfizer-BioNTech 6 months later (minimum interval 2 months) • Complete initial series with 1 dose Novavax at least 3 weeks after most recent dose, followed by 1 dose Moderna or Novavax or PfizerBioNTech 6 months later (minimum interval 2 months)
<p>Completed initial 3-dose vaccination series with:</p> <ul style="list-style-type: none"> • 3 or more doses Moderna or 3 or more doses PfizerBioNTech • 2 or more doses Novavax 	<ul style="list-style-type: none"> • 2 doses Moderna or Novavax or PfizerBioNTech 6 months apart (minimum interval 2 months). Administer dose 1 at least 8 weeks after the most recent dose • 2 doses Moderna or Novavax or Pfizer-BioNTech 6 months apart (minimum interval 2 months). Administer dose 1 at least 8 weeks after the most recent dose

Age 19 years or older

Vaccination Status	Recommended Doses
Unvaccinated	<ul style="list-style-type: none"> • 4 doses (3-dose initial series of Moderna at 0, 4 weeks, and at least 4 weeks after dose 2, followed by 1 dose Moderna or Novavax or Pfizer-BioNTech 6 months later [minimum interval 2 months]) • 4 doses (3-dose initial series of Pfizer-BioNTech at 0, 3 weeks, and at least 4 weeks after dose 2, followed by 1 dose Moderna or Novavax or Pfizer-BioNTech 6 months later [minimum interval 2 months]) • 3 doses (2-dose initial series Novavax at 0, 3 weeks, followed by 1 dose Moderna or Novavax or Pfizer-BioNTech 6 months later [minimum interval 2 months])
Incomplete initial vaccination series with: <ul style="list-style-type: none"> • 1 dose Moderna • 2 doses Moderna • 1 dose Pfizer-BioNTech • 2 doses Pfizer-BioNTech • 1 dose Novavax 	<ul style="list-style-type: none"> • Complete initial series with 2 doses Moderna at least 4 weeks apart (administer dose 1 4 weeks after most recent dose), followed by 1 dose Moderna or Novavax or Pfizer-BioNTech 6 months later (minimum interval 2 months) • Complete initial series with 1 dose Moderna at least 4 weeks after most recent dose, followed by 1 dose Moderna or Novavax or Pfizer-BioNTech 6 months later (minimum interval 2 months) • Complete initial series with 2 doses Pfizer-BioNTech at least 4 weeks apart (administer dose 1 3 weeks after most recent dose), followed by 1 dose Moderna or Novavax or Pfizer-BioNTech 6 months later (minimum interval 2 months) • Complete initial series with 1 dose Pfizer-BioNTech at least 4 weeks after most recent dose, followed by 1 dose Moderna or Novavax or Pfizer-BioNTech 6 months later (minimum interval 2 months) • Complete initial series with 1 dose Novavax at least 3 weeks after most recent dose, followed by 1 dose Moderna or Novavax or

	Pfizer-BioNTech 6 months later (minimum interval 2 months)
Completed the initial vaccination series with: <ul style="list-style-type: none"> • 3 or more doses Moderna or 3 or more doses Pfizer-BioNTech • 2 or more doses Novavax 	<ul style="list-style-type: none"> • 2 doses Moderna or Novavax or Pfizer-BioNTech 6 months apart (minimum interval 2 months). Administer dose 1 at least 8 weeks after the most recent dose • 2 doses Moderna or Novavax or Pfizer-BioNTech 6 months apart (minimum interval 2 months). Administer dose 1 at least 8 weeks after the most recent dose

Immunocompromised persons should discuss with their health care provider (shared clinical decision making) the need for additional doses of COVID-19 vaccine at least 2 months after the most recent dose. People can self-attest to being moderately or severely immunocompromised and receive additional COVID-19 vaccine doses.