



WOMEN

Symptoms

The majority of chlamydial infections in women do not cause any symptoms. You can get chlamydia in the cervix (opening to the womb), rectum, or throat. If you do have symptoms, you might notice:

- An unusual discharge from your vagina.
- Burning when you urinate.
- Discomfort or bleeding when you have sex.
- If the infection spreads, you might get lower abdominal pain, pain during sex, nausea, or fever.

Testing

You should be tested for chlamydia at least once a year if you are:

- Younger than 25 and you're having sex.
- 25 and older and you're having sex with more than one partner.
- 25 and older and you have a new sex partner.
- Pregnant.

Without Treatment

- If untreated, chlamydia can spread into the uterus or fallopian tubes and cause pelvic inflammatory disease (PID), a serious infection of the reproductive organs.
- PID can cause damage in your fallopian tubes. This damage may leave you unable to get pregnant or lead to an ectopic pregnancy (pregnancy outside the uterus).
- PID may also cause chronic pain in your pelvic area.
- If you have untreated chlamydia, you could pass the infection to your baby when giving birth. Chlamydia can cause serious health problems for babies.



MEN

Symptoms

The majority of chlamydial infections in men do not cause any symptoms. You can get chlamydia in the urethra (inside the penis), rectum, or throat. If you do have symptoms, you might notice:

- A discharge from your penis.
- Burning when you urinate.
- Burning or itching around the opening of your penis.

Testing

See a doctor if your partner has chlamydia or symptoms that might be chlamydia. Gay, bisexual, and men who have sex with men should also be tested for chlamydia.

Without Treatment

Chlamydia rarely causes long-term health problems in men. You may get an infection in the tube that carries sperm from the testes. This infection can cause pain and fever. In rare cases, this infection may prevent you from fathering children.

MEN AND WOMEN

See a doctor if your partner has chlamydia or symptoms that might be chlamydia.



Scan the Code
to find STI testing
centers near you.

Or, go to whitmancountypublichealth.org under Community Health and select the Sexually Transmitted Infections page.

**#GETTESTED
& WEAR PROTECTION**

THE FACTS

Many people who have chlamydia don't know it, and women often have no symptoms at all.

You can pass chlamydia to others without knowing it. Gonorrhea can be cured with the right treatment.

If you do not treat chlamydia, it can lead to serious health problems.

How Can I Lower My Risk For Chlamydia?

- The surest way to prevent chlamydia is not to have sex or to have sex only with someone who's not infected and who has sex only with you.
- Condoms can reduce your risk of getting chlamydia if used correctly every single time you have sex.
- Washing the genitals, urinating, or douching after sex will not prevent any STD.

How Does Someone Get Chlamydia?

- You can get chlamydia by having sex with someone who has it.
- "Having sex" means having anal, oral, or vaginal sex.
- If you are a pregnant woman with chlamydia, you can pass the infection to your baby.

How Can I Find Out If I Have Chlamydia?

- Ask a doctor to give you a test for chlamydia. The test is easy and painless.

If I Have Chlamydia, What Does That Mean For My Partner?

- Your partner may have chlamydia, too.
- Tell your recent sex partners so that they can get tested and treated.
- Avoid having sex until seven days after you've both started your treatment, so you don't re-infect each other.

How Is Chlamydia Treated?

- Chlamydia can be treated and cured with antibiotics.
- Finish all of the medicine to be sure you are cured. Do not share your medicine with anyone. You need all of it.
- If you still have symptoms after treatment, go back to see the doctor.
- You should get tested again about three months after you finish your treatment. This is especially important if you are unsure if your partner was also treated.

Can I Get Chlamydia Again After I've Been Treated?

- Yes, you can get chlamydia again. You can get it from an untreated partner or a new partner.

**#GETTESTED
& WEAR PROTECTION**



Whitman County
Public Health

PULLMAN
1205 SE Pro Mall Blvd #203
Pullman, WA 99163
509.332.6752

COLFAX
310 N Main Street #108
Colfax, WA 99111
509.397.6280